

Análisis di Enfermedad sexualmente transmití (SOA)

Instrukshon pa kolekshon di urina

1



Usa e pochi di urina ku bo a risibí di laboratorio.

2



Hasi uso di e promé urina di dia, òf sòru pa bo tin mínimo 1 ora sin uriná.

3



Fangu e promé stral di urina mes ora den e pochi. Yena e pochi ku mínimo 10 ML/CC; esaki ta marká riba e pochi. Yena e pochi mas tantu posibel. Kontinuá di urina den e wc. Kontrolá si e pochi di urina ta bon será.

4



Skibi e orario ku a kolekshoná e urina, riba e pochi.

5



Entrega e pochi ku urina denter di 2 ora, na un prekpòst di ADC den bo serkania.

Analysis of Sexually Transmitted Infections (STIs)

Instructions for collecting urine

1



Use the urine container provided by the laboratory.

2



Use the first morning urine, or ensure that you have not urinated for at least 1 hour before collecting the sample.

3



Urinate the first stream directly into the container. Fill the container with at least 10 ML/CC; the amount can be read on the container. Fill the container as much as possible. Then continue urinating into the toilet. Ensure that the container is properly sealed.

4



Write the time of urine collection on the container.

5



Submit the urine container within 2 hours to a nearby ADC blood collection center.

Analyse van Seksueel Overdraagbare Aandoeningen (SOA)

Instructies voor het verzamelen van urine

1



Gebruik het urinepotje dat u van het laboratorium heeft ontvangen.

2



Gebruik de eerste ochtendurine, óf zorg ervoor dat u minimaal 1 uur niet heeft geplast voordat u het monster verzamelt.

3



Plas het eerste straal urine direct in het potje. Vul het potje met minimaal 10 ML/CC; de hoeveelheid is af te lezen op het potje. Vul het potje zoveel mogelijk. Plas daarna verder uit in het toilet. Controleer of het potje goed is afgesloten.

4



Noteer het tijdstip van de urinemonsterafname op het potje.

5



Lever het potje met urine binnen 2 uur in bij een ADC prikpost bij u in de buurt.